# Native warm-season grass species and cultivars

### Patrick Keyser for Progressive Forage Grower

Recent studies indicate that native grasses can be a useful "tool" for forage producers, providing a good complement to tall fescue or orchardgrass and a low-input alternative to bermudagrass. There are five species typically considered for forage production: switchgrass, big bluestem, little bluestem, indiangrass and eastern gamagrass. While all have value for forage production, benefits and site adaptations vary by species and their more widely available cultivars.

### Switchgrass

This may be the most familiar of the native grasses to producers because of the attention it has received as a potential bioenergy crop. Switchgrass is very productive, typically yielding more than 5 tons per acre. In a recent University of Tennessee trial, single harvests of switchgrass yielded 3.5 and 5.5 tons per acre at early boot (late May) and early seed head (mid-June to late June) stages, respectively.

There are two basic types of switchgrass: upland and lowland. Lowland cultivars – Alamo and Kanlow are the two most common and are widely available – are taller and more stemmy, yielding about 25 to 50 percent more than many upland cultivars. Alamo and Kanlow are very similar, and the main distinction in terms of choosing which to plant is latitude. Growers from Kentucky north should consider the more cold-hardy Kanlow. Growers from Tennessee south will be better off with Alamo. Lowland cultivars, as their name implies, are adapted to very wet sites, producing acceptable yields where other commonproduction grasses will not survive.

There are many upland cultivars of switchgrass, but the highestyielding and best candidate for forage production is Cave-in-Rock. Like other upland switchgrasses, it has finer leaves and is less stemmy than lowland cultivars. Yields typically will be 10 to 20 percent lower than that produced by lowland cultivars.

In terms of forage quality, switchgrass can become stemmy, especially the lowland cultivars, but is readily grazed by cattle and produces summer gains of 1.5 to 2 pounds per day on steers. Carrying capacities (at 60 units nitrogen per acre) exceed 2,000 pounds per acre during May and June. Together, these attributes may make switchgrass the best option among the natives for overall forage production. Rates of gain are at a desirable level for backgrounding and heifer development, and the abundant growth allows for maintaining large numbers of cow-calf pairs during summer droughts. Indeed, switchgrass may be the most droughttolerant of the natives. Economic analyses conducted at the University of Tennessee indicate it provides the best returns and cheapest gains among native grasses.

### **Big bluestem**

Among the natives, big bluestem is the most preferred by cattle. Yields are not as high as switchgrass with annual production (one to two cuts per year) at about 4 tons per acre. This species has fine, leafy foliage, especially OZ-70 and Rountree cultivars. These two cultivars do well from Tennessee north with Earl (a Texas-origin cultivar) being better adapted to conditions in the Deep South. There are a number of very desirable local ecotypes being produced by seed growers, including Mammoth, Karst and Prairie. Kaw is an Oklahoma-origin cultivar that also does well in the eastern U.S.

Big bluestem produces excellent forage with steers gaining 2 to 2.5 pounds per day throughout the

summer. Carrying capacity for big bluestem is only about two-thirds of that produced by lowland switchgrass (1,000 pounds per acre during May and June). Big bluestem can grow on a wide variety of sites but does not do as well on wet sites as switchgrass. It does best on moderately to well-drained sites. It grows well in blends with indiangrass and little bluestem, and such blends can be managed compatibly for pasture or hay. For producers interested in heifer development or backgrounding steers, it is likely the best option because of the exceptional gains it can produce. It also would be the best choice for producing hay for the cash market.

### Little bluestem

Little bluestem produces

less forage than the other species mentioned here. However, it has the advantage of doing well on particularly poor sites and, compared to other natives, is easily established. Therefore, it also can be a good complement in blends with big bluestem and indiangrass by filling gaps in the stand where soil is poorer or establishment was not as successful. The most widely available cultivar is Aldous, although there are a number of other options available, including some desirable local ecotypes. On dry ridges or other sites with thin or degraded soil, a blend of little bluestem and indiangrass will be the best option for forage production.

Continued on page 34



## IT'S TIME TO CALL IN THE SPECIAL FORCES

The battle is on, and your silage deserves a higher level of protection. Unless you take appropriate action, yeasts, molds and other spoilage microbes can waste valuable feed or produce damaging toxins.

Order up Biotal<sup>®</sup> forage inoculants and ensure the silage you put in is the silage you get out. With an





## Make the call to your Lallemand Animal Nutrition representative to have Biotal forage inoculants protect your silage.

BIOTAL

### www.Biotal.com

 LALLEMAND ANIMAL NUTRITION
 www.lallemandani

 Tel: 414 464 6440 Email: LAN\_NA@lallemand.com
 @2015. Biolal is a registered trademark of Lallemand Animal Nutrition. Not all products are available in all markets nor are all claims allowed in all regions.



## Native warm-season grass species and cultivars, cont'd from page 33

#### Indiangrass

Compared to big bluestem, indiangrass is somewhat more productive (about 5 tons per acre annual production based on one to two cuts per year), its leaves are not as fine, and it is slightly more stemmy. Indiangrass is an excellent forage, being only slightly less preferred by cattle than big bluestem. Like little bluestem, it is one of the easier natives to establish. The better-known cultivars that do well in the eastern U.S. are Rumsey and Osage. Americus, a new release based on southern genetics, is well adapted to the Deep South. Additionally, there are several relatively new local ecotypes on the market that may provide better adaptation based on their origin; they include Boone (Kentucky origin), Prairie (Indiana origin) and VA (Virginia origin).

Typically, steers do about as well on this species as they do on big bluestem, with gains exceeding 2 pounds per day. Carrying capacity is somewhat higher than that of big bluestem though, at about 1,200 pounds per acre at peak early summer growth. Indiangrass is less tolerant of wet sites than big bluestem. It can grow on a wide variety of soils, being intermediate between big and little bluestem in terms of its tolerance of poor soils. Like big bluestem, it is an excellent choice for backgrounding steers or heifer development and grows well in blends with that species.

### Eastern gamagrass

In terms of yield, carrying capacity and drought tolerance, eastern gamagrass is comparable to switchgrass. However, it starts growing earlier in the spring and sustains production in late summer better than the others and, unlike the other natives, is very responsive to nitrogen. There

Does your fertilizer match your spreader performance?

The spreadability of fertilizer is affected by many things, not the least of which is fertilizer quality. High spreading distance and even spreading patterns are important to the success of your crops, but using low-quality fertilizers can greatly limit high-performance spreader effectiveness.

Propelling with high acceleration and force puts severe physical strain on fertilizer granules. Ammonium nitrate, due to a higher bulk density and lower nitrogen concentration, offers more homogeneous spreading over the field than urea. Spreading nitrogen fertilizer precisely increases nutrient availability to crops and reduces environmental losses.

Yara's nitrate-based fertilizers are pure nutrients. They offer nutrient efficiency, environmental sustainability and greater yield. Pure Nutrients. Scan for more information on Yara's crop nutrition products.





Patrick Keyser Director Center for Native Grasslands Management University of Tennessee pkeyser@utk.edu

#### www.progressiveforage.com

are three relatively common cultivars: Pete, Iuka and Highlander. The first two are similar and have finer leaves and lower yields (about 25 percent less) than Highlander. All three cultivars do well throughout the eastern U.S.

From a quality standpoint, studies at the University of Tennessee indicate gains (steer-weight basis) of about 1.4 pounds per day are reasonable. In terms of carrying capacity, a University of Tennessee trial in which no nitrogen was applied supported stocking of more than 2,000 pounds per acre with Highlander. Like lowland switchgrass, eastern gamagrass can tolerate very wet sites. On the other hand, it is not as tolerant of poor sites as the other natives discussed here. Because of the modest rate of gain for this species, it would not be as desirable for stockers or heifer development as the other species. However, because of its high carrying capacity, eastern gamagrass can be a very useful grass for providing forage during prolonged droughts. Producers with access to inexpensive forms of nitrogen (poultry litter, dairy effluent, municipal sludge) could experience considerably higher gains and carrying capacities given this species' ability to respond to fertility.

Consideration of the site adaptations, available cultivars and attributes of these species can allow producers to identify options among native grasses that may be best suited to their operations.

Additional information is available at the UT Extension publications website by entering the term "native warm-season grass" into the search engine (utextension. tennessee.edu/publications) or visit the website for the Center for Native Grasslands Management (nativegrasses.utk.edu). **FG** 



VARA.